


SLC18A2 Antibody / VMAT2 [clone SLC18A2/7983] (V4123)


Catalog No.	Formulation	Size
V4123-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4123-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4123SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2, kappa
Clone Name	SLC18A2/7983
Purity	Protein A/G affinity
UniProt	Q05940
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This SLC18A2 antibody is available for research use only.



IHC staining of FFPE human brain tissue with SLC18A2 antibody (clone SLC18A2/7983). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate tissue with SLC18A2 antibody (clone SLC18A2/7983). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free SLC18A2 antibody (clone SLC18A2/7983) as confirmation of integrity and purity.

Description

Neurotransmission depends on the regulated exocytotic release of chemical transmitter molecules. This requires the packaging of these substances into the specialized secretory vesicles of neurons and neuroendocrine cells, a process mediated by specific vesicular transporters. The family of genes encoding the vesicular transporters of monoamines (VMAT 1 and VMAT 2) and acetylcholine (VACht) have been cloned and functionally characterized. The sequence of these integral membrane proteins predicts twelve transmembrane domains and weak homology to a class of bacterial antibiotic resistance proteins. The vesicular transport of neurotransmitter molecules has been shown to be an active ATP- and proton dependent transport mechanism.

Application Notes

Optimal dilution of the SLC18A2 antibody should be determined by the researcher.

Immunogen

A recombinant fragment of human SLC18A2 protein was used as the immunogen for the SLC18A2 antibody.

Storage

Aliquot the SLC18A2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.