


SM-MHC Antibody [clone SMMS-1] (V2748)

Catalog No.	Formulation	Size
V2748-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2748-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2748SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2748IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


 Citations (11)

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SMMS-1
Purity	Protein G affinity chromatography
UniProt	P35749
Localization	Cytoplasmic
Applications	Flow Cytometry : 1-2ug/million cells in 0.1ml Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT (1) Prediluted IHC Only Format : incubate for 30 min at RT (2)
Limitations	This SM-MHC antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human Leiomyosarcoma stained with SM-MHC antibody (clone SMMS-1).



IHC testing of FFPE human breast carcinoma with SM-MHC antibody (clone SMMS-1).

Description

Smooth muscle myosin heavy chain (SM-MHC) is a cytoplasmic structural protein, which is a major component of the contractile apparatus in smooth muscle cells. Expression of smooth muscle myosin is developmentally regulated, appearing early in smooth muscle development, and is specific for smooth muscle development. Two isoforms of smooth muscle myosin heavy chain have been identified, designated MHC-1 and MHC-2. The antibody may be useful for the study of breast tumors as the presence of an intact layer of myoepithelial cells is an important feature, which may distinguish benign breast lesions and carcinoma in situ from invasive tumors.

Application Notes

The stated application concentrations are suggested starting amounts. Optimal dilution of the SM-MHC antibody should be determined by the researcher.

1. Staining of formalin/paraffin tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Human uterus extract was used as the immunogen for the SM-MHC antibody.

Storage

Store the SM-MHC antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).