


SMAD2 Antibody (R35875)


| Catalog No. | Formulation | Size |
|--------------|---|--------|
| R35875-100UG | 0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide | 100 ug |

[Bulk quote request](#)

| | |
|-----------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human, Mouse |
| Predicted Reactivity | Cow |
| Format | Antigen affinity purified |
| Host | Goat |
| Clonality | Polyclonal (goat origin) |
| Isotype | Goat Ig |
| Purity | Antigen affinity |
| Gene ID | 4087 |
| Applications | Western Blot : 2-4ug/ml ELISA (peptide) LOD : 1:4000 |
| Limitations | This SMAD2 antibody is available for research use only. |



Western blot testing of HeLa lysate with SMAD2 antibody at 2ug/ml. The 467 AA and 437 AA isoforms may be detected with this Ab.



Western blot testing of mouse testis lysate with SMAD2 antibody at 2ug/ml.

Description

Additional name(s) for this target protein: Mothers against decapentaplegic homolog 2, MADH2

Application Notes

Optimal dilution of the SMAD2 antibody should be determined by the researcher.

Immunogen

Amino acids SEIWGLSTPNTIDC were used as the immunogen for this SMAD2 antibody.

Storage

Aliquot and store the SMAD2 antibody at -20oC.