
# SMMHC Antibody / MYH11 [clone MYH11/7610R] (V4416)

| Catalog No. | Formulation | Size |
|---|---|---|
| V4416-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4416-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V4416SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Recombinant RABBIT MONOCLONAL

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Purified |
| **Host** | Rabbit |
| **Clonality** | Recombinant Rabbit Monoclonal |
| **Isotype** | Rabbit IgG, kappa |
| **Clone Name** | MYH11/7610R |
| **Purity** | Protein A/G affinity |
| **UniProt** | P35749 |
| **Localization** | Cytoplasm |
| **Applications** | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT |
| **Limitations** | This SMMHC antibody is available for research use only. |



IHC staining of FFPE human smooth muscle tissue with SMMHC antibody (clone MYH11/7610R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## Description

Smooth Muscle Myosin, heavy chain (SMM-HC) is a cytoplasmic structural protein that is a major component of the contractile apparatus of the smooth muscle cells. Expression of smooth muscle myosin is developmentally regulated,

appearing early and is specific for smooth muscle development. SM-MHC stains the intact myoepithelial cell (MEC) layers present in benign and in situ malignant breast and bronchioloalveolar lesions and is therefore very helpful in distinguishing between benign and malignant tumors. The antibody reacts with smooth muscle cells and myoepithelial cells, but not with myofibroblasts. It is very helpful in distinguishing between benign sclerosing breast lesions and infiltrating carcinomas in difficult cases since it strongly stains the myoepithelial layer in the benign lesions while it is negative in the infiltrating carcinomas.

Visit our SMMHC Antibody page to discover additional antibodies against this Smooth Muscle Differentiation Marker, including reagents for studies of vascular biology, contractility, and smooth muscle pathology.

## Application Notes

Optimal dilution of the SMMHC antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 1-200) from the human protein was used as the immunogen for the SMMHC antibody.

## Storage

Aliquot the SMMHC antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.


Copyright © NSJ Bioreagents. All rights reserved.