



SPARC Antibody / Osteonectin [clone OSTN/3304] (V9733)

Catalog No.	Formulation	Size
V9733-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9733-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9733SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	OSTN/3304
Purity	Protein A/G affinity
UniProt	P09486
Localization	Secreted
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This SPARC antibody is available for research use only.





IHC staining of FFPE human tonsil tissue with SPARC antibody (clone OSTN/3304).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free SPARC antibody (clone OSTN/3304) as confirmation of integrity and purity.

Description

SPARC (for secreted protein acidic and rich in cysteine) is a phosphorylated, acidic, glycine-rich glycoprotein that is secreted by endothelial cells and is present in large amounts in the parietal endoderm of mouse embryos and in human placenta. It is identical to osteonectin, a protein important to bone calcification that is highly conserved between species. SPARC, which can be selectively expressed by the endothelium in response to certain types of injury, induces rounding in adherent endothelial cells in vitro. It regulates endothelial barrier function through F-Actin-dependent changes in cell shape, coincident with the appearance of intercellular gaps, which provide a paracellular pathway for extravasation of macromolecules.

Application Notes

Optimal dilution of the SPARC antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-200 was used as the immunogen for the SPARC antibody.

Storage

Aliquot the SPARC antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.