


SPTLC1 Antibody / Serine palmitoyltransferase 1 (R30870)


Catalog No.	Formulation	Size
R30870	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days	
Species Reactivity	Human, Mouse, Rat	
Format	Antigen affinity purified	
Clonality	Polyclonal (rabbit origin)	
Isotype	Rabbit IgG	
Purity	Antigen affinity	
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal	
UniProt	O15269	
Localization	Cytoplasmic	
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml	
Limitations	This SPTLC1 antibody is available for research use only.	



Western blot testing of 1) human SiHa, 2) human U-251, 3) rat brain and 4) mouse brain tissue lysate with SPTLC1 antibody. Expected molecular weight ~53 kDa.



IHC staining of FFPE human breast cancer tissue with SPTLC1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

Serine palmitoyltransferase, long chain base subunit 1, also known as SPT1, LCB1, is a protein which in humans is encoded by the SPTLC1 gene. Dawkins et al.(2001)noted that the gene maps to chromosome 9q22.1-q22.3. Serine palmitoyltransferase, which consists of two different subunits, is the initial enzyme in sphingolipid biosynthesis. It converts L-serine and palmitoyl CoA to 3-oxosphinganine with pyridoxal 5'-phosphate as a cofactor. The product of this gene is the long chain base subunit 1 of serine palmitoyltransferase. Mutations in this gene were identified in patients with hereditary sensory neuropathy type 1. Alternatively spliced variants encoding different isoforms have been identified.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the SPTLC1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the middle region of human SPTLC1 (FYGTFDVHLDLEDRLAK) was used as the immunogen for this SPTLC1 antibody.

Storage

After reconstitution, the SPTLC1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.