

SRC-1 Antibody / NCOA1 (F50522)

Catalog No.	Formulation	Size
F50522-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F50522-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Pig
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
UniProt	Q15788
Applications	Western Blot : 1:1000 IHC (Paraffin) : 1:10-1:50
Limitations	This SRC-1 antibody is available for research use only.

250
130
95
72
55

SRC-1 antibody western blot analysis in CEM lysate.



IHC analysis of FFPE human colon carcinoma tissue stained with SRC-1 antibody

Description

SRC-1/NCOA1 acts as a transcriptional coactivator for steroid and nuclear hormone receptors. It is a member of the p160/steroid receptor coactivator (SRC) family and like other family members has histone acetyltransferase activity and contains a nuclear localization signal, as well as bHLH and PAS domains. This protein binds nuclear receptors directly and stimulates the transcriptional activities in a hormone-dependent fashion.

Application Notes

Titration of the SRC-1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 1338-1367 from the human protein was used as the immunogen for this SRC-1 antibody.

Storage

Aliquot the SRC-1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.