


STAT1 Antibody (R32095)


Catalog No.	Formulation	Size
R32095	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P42224
Localization	Cytoplasmic
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This STAT1 antibody is available for research use only.




Western blot testing of rat 1) testis, 2) brain, 3) liver, and human 4) placenta, 5) MCF7, and 6) SW620 lysate with STAT1 antibody. Predicted/observed molecular weight: ~91/84 kDa (alpha/beta).



IHC testing of FFPE human breast cancer with STAT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE mouse intestine with STAT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat intestine with STAT1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

Description

The crystal structure of the DNA complex of a 67-kD core fragment of the STAT1 homodimer was determined, lacking only the N-domain and the C-terminal transcriptional activation domain, at 2.9-angstrom resolution. Phosphorylation of Signal Transducer and Activator of transcription 1 (STAT 1) was also decreased in rheumatoid arthritis lymphocytes. The transcription factor signal transducer and activator of transcription-1 (STAT1) plays a key role in immunity against mycobacterial and viral infections. Activation of the signal transducers and activators of transcription (STAT) pathway is important in fibroblast growth factor (FGF) modulation of chondrocyte proliferation and endochondral bone formation during embryogenesis.

Application Notes

Optimal dilution of the STAT1 antibody should be determined by the researcher.

Immunogen

Amino acids KILENAQRFNQAQSGNIQSTVMLDKQKELD of human STAT1 were used as the immunogen for the STAT1 antibody.

Storage

After reconstitution, the STAT1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.