


## STXBP2 Antibody / UNC18-2 (R32239)


Catalog No.	Formulation	Size
R32239	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
<b>UniProt</b>	Q15833
<b>Applications</b>	Western Blot : 0.1-0.5ug/ml Immunofluorescence (FFPE) : 2-4ug/ml Flow Cytometry : 1-3ug/million cells
<b>Limitations</b>	This STXBP2 antibody is available for research use only.



Immunofluorescent staining of FFPE human SK-OC-3 cells with STXBP2 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human U937 cells with STXBP2 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= STXBP2 antibody.



Western blot testing of 1) rat brain, 2) mouse brain, 3) human HepG2 and 4) human HeLa lysate with STXBP2 antibody. Expected molecular weight ~66 kDa.

## Description

Syntaxin-binding protein 2, also known as UNC18B and UNC18-2, is a protein that in humans is encoded by the STXBP2 gene. The STXBP2 gene is mapped to human chromosome 19p13.3-p13.2 and to the proximal arm of mouse chromosome 8. And this gene encodes a member of the STXBP/unc-18/SEC1 family. The encoded protein is involved in intracellular trafficking, control of SNARE (soluble NSF attachment protein receptor) complex assembly, and the release of cytotoxic granules by natural killer cells. Additionally, UNC18B is expressed predominantly as a 2.4-kb message in placenta, lung, liver, kidney, and pancreas, as well as in peripheral blood lymphocytes. Mutations in this gene are associated with familial hemophagocytic lymphohistiocytosis. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene.

## Application Notes

Optimal dilution of the STXBP2 antibody should be determined by the researcher.

## Immunogen

Amino acids QEYPAIRYRKGPEDTAQLAHAVLAKLNAFKAD of human STXBP2 were used as the immunogen for the STXBP2 antibody.

## Storage

After reconstitution, the STXBP2 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.