

T-Cell Intracellular Antigen-1 Antibody / TIA-1 [clone rTIA1/1353] (V4053)

Catalog No.	Formulation	Size
V4053-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4053-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4053SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rTIA1/1353
Purity	Protein A affinity
UniProt	P31483
Localization	Cytoplasm, nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This T-Cell Intracellular Antigen-1 antibody is available for research use only.



Description

The T cell intracellular antigen 1 (TIA 1) is a 17 kDa cytoplasmic granule associated protein also designated as GMP 17, for granule membrane protein of 17 kDa. The GMP 17/TIA 1 molecule is expressed in cells possessing cytolytic potential

and could be involved in the signaling cascade of Fas (CD95) mediated apoptosis. Within hematopoietic cell lines, the rTIA1/1353 monoclonal antibody (mAb) reacts with about 90% of CD16+, 50-60% of CD8+, and less than 10% of CD4+ normal peripheral blood lymphocytes. It reacts with almost all monocytes and granulocytes. This antibody also reacts with CD4+ activated T cell clones, activated NK cell clones, and Con activated thymocytes, but not with B lymphocytes or B cell lines. TIA1 antibody labels cytotoxic T cells and natural killer cells (NK cells). It is also expressed in T-cell lymphoma, large granular lymphocyte (LGL) leukemia and hairy cell leukemia. TIA1 expression in T-cell malignancies may help in differentiating LGL leukemia (high expression) from T-cell lymphocytosis and other T-cell diseases (low expression). TIA1 may also be used to label tumor-infiltrating lymphocytes in the study of immune response to malignancies.

Application Notes

Optimal dilution of the T-Cell Intracellular Antigen-1 antibody should be determined by the researcher.

Immunogen

Recombinant full-length human protein was used as the immunogen for the T-Cell Intracellular Antigen-1 antibody.

Storage

Aliquot the T-Cell Intracellular Antigen-1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.