


TACR1 Antibody (R32031)

Catalog No.	Formulation	Size
R32031	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P25103
Applications	Western Blot : 0.1-0.5ug/ml
Limitations	This TACR1 antibody is available for research use only.



Western blot testing of 1) human placenta, 2) rat lung, 3) rat brain, 4) U87 and 5) A431 lysate with TACR1 antibody. Predicted molecular weight ~46 kDa but routinely observed at ~58 kDa.

Description

The tachykinin receptor 1 (TACR1), also known as neurokinin 1 receptor (NK1R) or substance P receptor (SPR), is a G protein coupled receptor found in the central nervous system and peripheral nervous system. The endogenous ligand for this receptor is Substance P, although it has some affinity for other tachykinins. The protein is the product of the TACR1 gene. Tachykinin receptor 1 consists of 407 amino acid residues, and it has a molecular weight of 58,000D. Tachykinin receptor 1, as well as the other tachykinin receptors, is made of seven hydrophobic transmembrane (TM) domains with three extracellular and three intracellular loops, an amino-terminus and a cytoplasmic carboxy-terminus. The loops have functional sites, including two cysteines amino acids for a disulfide bridge, Asp-Arg-Tyr, which is responsible for

association with arrestin and, Lys/Arg-Lys/Arg-X-X-Lys/Arg, which interacts with G-proteins. The tachykinin receptor 1 can be found in both the central and peripheral nervous system. It is present in neurons, brainstem, vascular endothelial cells, muscle, gastrointestinal tracts, genitourinary tract, pulmonary tissue, thyroid gland and different types of immune cells. The binding of SP to the tachykinin receptor 1 has been associated with the transmission of stress signals and pain, the contraction of smooth muscles and inflammation. Tachykinin receptor 1 antagonists have also been studied in migraine, emesis and psychiatric disorders. In fact, aprepitant has been proved effective in a number of pathophysiological models of anxiety and depression. Other diseases in which the tachykinin receptor 1 system is involved include asthma, rheumatoid arthritis and gastrointestinal disorders.

Application Notes

Optimal dilution of the TACR1 antibody should be determined by the researcher.

Immunogen

Amino acids FHIFFLLPYINPDLYLKKFIQQVYLAIM of human TACR1 were used as the immunogen for the TACR1 antibody.

Storage

After reconstitution, the TACR1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.