


## TAFI Antibody / CPB2 (R32483)

| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| R32483      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

**Bulk quote request**

|                           |   |
|---------------------------|---|
| <b>Availability</b>       | 1-3 business days   |
| <b>Species Reactivity</b> | Mouse   |
| <b>Format</b>             | Antigen affinity purified                                     |
| <b>Clonality</b>          | Polyclonal (rabbit origin)                                    |
| <b>Isotype</b>            | Rabbit IgG  |
| <b>Purity</b>             | Antigen affinity  |
| <b>Buffer</b>             | Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide |
| <b>UniProt</b>            | Q9JHH6  |
| <b>Applications</b>       | Western Blot : 0.5-1ug/ml                                     |
| <b>Limitations</b>        | This TAFI antibody is available for research use only.        |



Western blot testing of mouse liver lysate with TAFI antibody at 0.5ug/ml. Predicted molecular weight: ~48 kDa but routinely observed at 50-60 kDa.

## Description

Carboxypeptidase B2 (CPB2), also known as carboxypeptidase U (CPU), plasma carboxypeptidase B (pCPB) or thrombin-activatable fibrinolysis inhibitor (TAFI), is an enzyme that, in humans, is encoded by the gene CPB2. CPB2 is synthesized by the liver and circulates in the plasma as a plasminogen-bound zymogen. When it is activated by proteolysis at residue Arg92 by the thrombin/thrombomodulin complex, CPB2 exhibits carboxypeptidase activity. Activated CPB2 reduces fibrinolysis by removing the fibrin C-terminal residues that are important for the binding and activation of plasminogen.

## Application Notes

Optimal dilution of the TAFI antibody should be determined by the researcher.

## **Immunogen**

Amino acids K165-D387 were used as the immunogen for the TAFI antibody.

## **Storage**

Prior to reconstitution, store at 4oC. After reconstitution, the TAFI antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.