


## TAK1 Antibody / MAP3K7 (F50911)

Catalog No.	Formulation	Size
F50911-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F50911-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Predicted Reactivity</b>	Bovine, Mouse, Rat
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Purified
<b>UniProt</b>	O43318
<b>Applications</b>	Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100
<b>Limitations</b>	This TAK1 antibody is available for research use only.



IHC analysis of FFPE human hepatocarcinoma tissue stained with the TAK1 antibody.



Western blot analysis of TAK1 antibody and HL-60 cell lysate. Predicted molecular weight: 64-69 kDa, routinely observed at 78-82 kDa.

## Description

TAK1 is a member of the serine/threonine protein kinase family. This kinase mediates the signaling transduction induced by TGF beta and morphogenetic protein (BMP), and controls a variety of cell functions including transcription regulation and apoptosis. In response to IL-1, this protein forms a kinase complex including TRAF6, MAP3K7P1/TAB1 and MAP3K7P2/TAB2; this complex is required for the activation of nuclear factor kappa B. This kinase can also activate MAPK8/JNK, MAP2K4/MKK4, and thus plays a role in the cell response to environmental stresses.

## Application Notes

Titration of the TAK1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 574-606 from the human protein was used as the immunogen for this TAK1 antibody.

## Storage

Aliquot the TAK1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.