


TARDBP Antibody / TAR DNA binding protein / TDP43 [clone TARDP/349] (V4706)


Catalog No.	Formulation	Size
V4706-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4706-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4706SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1
Clone Name	TARDP/349
Purity	Protein A/G affinity
UniProt	Q13148
Localization	Nucleus
Applications	ELISA (Order BSA-free Format For Coating) : Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This TARDBP antibody is available for research use only.



IHC staining of FFPE human prostate tissue with TARDBP antibody (clone TARDP/349).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free TARDBP antibody (clone TARDP/349) as confirmation of integrity and purity.

Description

TARDBP (TAR DNA binding protein), also known as TDP-43, is a nuclear protein that contains two RRM (RNA recognition motif) domains. Ubiquitously expressed with highest levels found in placenta, lung, pancreas, spleen and genital tract, TARDBP functions as a DNA-binding protein and specifically binds to the TAR DNA sequence motifs of HIV. Via this association with TAR motifs, TARDBP acts as a transcriptional repressor and inhibits HIV-1 transcription. TARDBP can also function as a negative regulator of splicing activity and is known to be involved in the splicing of CFTR (cystic fibrosis transmembrane receptor). In addition, TARDBP is a major component of ubiquitin-positive inclusion bodies that are prominent in many neurodegenerative diseases. This suggests that TARDBP may play a role in the development of neurodegenerative disorders. Due to alternative splicing events, various isoforms exist for TARDBP.

Application Notes

Optimal dilution of the TARDBP antibody should be determined by the researcher.

Immunogen

A recombinant fragment from the human protein (within amino acids 200-414) was used as the immunogen for the TARDBP antibody.

Storage

Aliquot the TARDBP antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.