


# TDRKH Antibody [clone PCRP-TDRKH-1H2] (V9602)


Catalog No.	Formulation	Size
V9602-100UG	0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9602-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9602SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	PCRP-TDRKH-1H2
Purity	Protein A/G affinity
UniProt	Q9Y2W6
Localization	Cytoplasm
Applications	Flow Cytometry: 1-2ug/million cells Immunofluorescence: 1-2ug/ml Western Blot: 1-2ug/ml Immunohistochemistry (FFPE): 1-2ug/ml
Limitations	This TDRKH antibody is available for research use only.




Immunofluorescent staining of PFA-fixed human HeLa cells using TDRKH antibody (green, clone PCRP-TDRKH-1H2) and phalloidin (red).




IHC staining of FFPE human ovarian carcinoma tissue with TDRKH antibody (clone PCRP-TDRKH-1H2) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



FACS staining of PFA-fixed human HeLa cells with TDRKH antibody (blue, clone PCRP-TDRKH-1H2), and unstained cells (red).



SDS-PAGE analysis of purified, BSA-free TDRKH antibody (clone PCRP-TDRKH-1H2) as confirmation of integrity and purity.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using TDRKH antibody (clone PCRP-TDRKH-1H2). These results demonstrate the foremost specificity of the PCRP-TDRKH-1H2 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

### **Description**

Participates in the primary piRNA biogenesis pathway and is required during spermatogenesis to repress transposable elements and prevent their mobilization, which is essential for the germline integrity. The piRNA metabolic process mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and govern the methylation and subsequent repression of transposons. Required for the final steps of primary piRNA biogenesis by participating in the processing of 31-37 nt intermediates into mature piRNAs. May act in pi-bodies and piP-bodies by transferring piRNA precursors or intermediates to or between these granules.

## **Application Notes**

Optimal dilution of the TDRKH antibody should be determined by the researcher.

### **Immunogen**

Recombinant full-length human TDRKH protein was used as the immunogen for the TDRKH antibody.

# **Storage** Aliquot the TDRKH antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.