

## TLR4 Antibody (Toll Like Receptor 4) [clone TLR4/230] (V2897)

Catalog No.	Formulation	Size
V2897-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2897-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2897SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a, kappa
<b>Clone Name</b>	TLR4/230
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	O00206
<b>Localization</b>	Cell surface
<b>Applications</b>	Functional Studies (order BSA/sodium Azide-free Format) :
<b>Limitations</b>	This TLR4 antibody is available for research use only.



### Description

This mAb reacts with human Toll-like receptor 2 (TLR4). It is a member of the Toll-like receptor (TLR) family, which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from *Drosophila* to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor has been implicated in signal transduction events induced by lipopolysaccharide (LPS) found in most gram-negative bacteria. Mutations in this gene have been associated with differences in LPS responsiveness. Multiple transcript variants encoding different isoforms have been found for this gene.

## Application Notes

Optimal dilution of the TLR4 antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.

## Immunogen

Recombinant human protein was used as the immunogen for the TLR4 antibody.

## Storage

Store the TLR4 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).