


TNF Receptor Antibody / TNFR1 / TNFRSF1A (RQ4614)


Catalog No.	Formulation	Size
RQ4614	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P19438
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml ELISA (Capture) : 1-5ug/ml (recombinant human protein)
Limitations	This TNF Receptor antibody is available for research use only.




IHC staining of FFPE mouse colon tissue with TNF Receptor antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse kidney tissue with TNF Receptor antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) HeLa and 2) A549 cell lysate with TNF Receptor antibody. Predicted molecular weight ~51 kDa.



Western blot testing of 1) rat brain, 2) rat lung, 3) mouse brain and 4) mouse lung tissue lysate with TNF Receptor antibody. Predicted molecular weight ~51 kDa.



Western blot testing of 1) human HeLa, 2) human K562, 3) human Caco-2, 4) rat liver, 5) mouse small intestine, 6) mouse liver and 7) mouse HEPA1-6 lysate with TNF Receptor antibody at 0.5ug/ml. Predicted molecular weight ~51 kDa.

Description

Tumor necrosis factor receptor superfamily member 1A (TNFRSF1A), also known as TNFR1, is a protein that in humans is encoded by the TNFRSF1A gene. The protein encoded by this gene is a member of the Tumor necrosis factor receptor superfamily, which also contains TNFRSF1B. The TNFR1 gene is mapped to 12pter-cen. It encodes a protein of 455 amino acids. And this receptor can activate the transcription factor NF- κ B, mediate apoptosis, and function as a regulator of inflammation.

Application Notes

Optimal dilution of the TNF Receptor antibody should be determined by the researcher.

Immunogen

Amino acids I22-T211 from the human protein were used as the immunogen for the TNF Receptor antibody.

Storage

After reconstitution, the TNF Receptor antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

