

# TNFAIP8 Antibody / Tumor necrosis factor alpha induced protein 8 [clone 30T22] (FY12872)

| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| FY12872     | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium | 100 ul |
|             | azide and 50% glycerol, 0.4-0.5mg/ml BSA                                  |        |

## Recombinant RABBIT MONOCLONAL

## **Bulk quote request**

| Availability       | 2-3 weeks   |
|--------------------|---|
| Species Reactivity | Human   |
| Format             | Liquid  |
| Clonality          | Recombinant Rabbit Monoclonal   |
| Isotype            | Rabbit IgG  |
| Clone Name         | 30T22   |
| Purity             | Affinity-chromatography   |
| Buffer             | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA. |
| UniProt            | O95379  |
| Applications       | Western Blot : 1:500-1:2000   |
| Limitations        | This TNFAIP8 antibody is available for research use only.   |

## **Description**

TNFAIP8 antibody detects Tumor necrosis factor alpha induced protein 8, encoded by the TNFAIP8 gene. This protein is part of the TNFAIP8 family, which includes several paralogs involved in apoptosis regulation, immune modulation, and cancer biology. TNFAIP8 has been described as an anti apoptotic factor that suppresses programmed cell death and influences cell survival pathways in response to cytokine signaling. Research using TNFAIP8 antibody has demonstrated that it contributes to immune evasion in tumors and enhances the survival of malignant cells under stress conditions, making it highly relevant in oncology studies.

TNFAIP8 is induced by TNF signaling and is involved in modulating NF kappa B activity. Studies indicate that it can inhibit caspase activation and reduce apoptosis, thereby promoting cell survival under stress conditions. By altering apoptosis thresholds, TNFAIP8 helps cancer cells persist in hostile microenvironments where other cells would undergo death. Detection with TNFAIP8 antibody has shown elevated expression in multiple cancers, including lung, colon, and ovarian carcinoma, where it contributes to chemoresistance and poor prognosis. Its regulation of immune signaling pathways

makes it a target of interest in both immunology and oncology, as it may serve as a biomarker for aggressiveness and therapeutic resistance.

Beyond its role in apoptosis, TNFAIP8 may participate in vesicle trafficking and autophagy. It has been implicated in lipid binding and membrane dynamics, linking it to cellular stress responses and nutrient sensing. Evidence suggests that TNFAIP8 can modulate PI3K AKT signaling, further highlighting its role in integrating survival and metabolic pathways. Researchers employing TNFAIP8 antibody can explore these broader functions and their relevance in pathological conditions, including inflammatory diseases where dysregulated cell survival exacerbates tissue injury.

TNFAIP8 antibody is commonly used in western blotting, immunohistochemistry, and ELISA. Western blotting identifies isoform expression and quantitative changes, while immunohistochemistry reveals expression patterns in tumor samples and immune tissues, offering insights into microenvironmental interactions. ELISA based assays extend applications to serum measurements in clinical samples, where circulating TNFAIP8 may serve as a biomarker for disease severity. Flow cytometry applications have also been developed, enabling detection in heterogeneous cell populations.

Research into tumor biology, inflammation, and therapeutic resistance benefits from reliable antibodies such as TNFAIP8 antibody. By providing validated reagents, NSJ Bioreagents supports discovery of the molecular functions and disease associations of Tumor necrosis factor alpha induced protein 8, ensuring researchers have consistent and reproducible tools to study this important factor.

## **Application Notes**

Optimal dilution of the TNFAIP8 antibody should be determined by the researcher.

### **Immunogen**

A synthesized peptide derived from human TNFAIP8 was used as the immunogen for the TNFAIP8 antibody.

#### **Storage**

Store the TNFAIP8 antibody at -20oC.