


## TOX-3 Antibody [clone TOX3/1124] (V2542)


Catalog No.	Formulation	Size
V2542-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2542-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2542SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	TOX3/1124
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	O15405
<b>Localization</b>	Cytoplasmic & nuclear
<b>Applications</b>	ELISA (order BSA/sodium Azide-free Format For Coating) : Western Blot : 1-2ug/ml
<b>Limitations</b>	This TOX-3 antibody is available for research use only.



Western blot analysis of 1) recombinant partial protein, 2) A549 and 3) A431 stained with TOX-3 antibody (clone TOX3/1124). Predicted molecular weight ~63 kDa.



SDS-PAGE Analysis of Purified, BSA-Free TOX-3 Antibody (clone TOX3/1124).  
Confirmation of Integrity and Purity of the Antibody.

## Description

It recognizes an ~63 kDa protein, which is identified as TOX3. It contains a high mobility group (HMG)-box, which regulates Ca<sup>2+</sup>-dependent transcription in neurons through interaction with the cAMP-response-element-binding protein (CREB). TOX3 appears to be associated with breast cancer susceptibility and is expressed downstream of a cytoprotective cascade together with CITED1, a transcriptional regulator that does not bind directly to DNA. TOX3 is predominantly expressed in the brain and forms homodimers. TOX3 overexpression protects neuronal cells from cell death caused by endoplasmic reticulum stress or BAX overexpression through the induction of anti-apoptotic transcripts and repression of pro-apoptotic transcripts.

## Application Notes

Optimal dilution of the TOX-3 antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

A recombinant fragment (139 amino acid residues around aa 200-400) from the human protein was used as the immunogen for the TOX-3 antibody.

## Storage

Store the TOX-3 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).