



# TRAF2 Antibody (R32205)

Catalog No.	Formulation	Size
R32205	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	Q12933
Localization	Cytoplasmic
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This TRAF2 antibody is available for research use only.





IHC testing of FFPE human tonsil with TRAF2 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

#### **Description**

TRAF2 (TNF Receptor-Associated Factor 2), also called TRAP, is a protein that in humans is encoded by the TRAF2 gene. The protein encoded by this gene is a member of the TNF receptor (TNFR) associated factor (TRAF) protein family. It is a common signal transducer for TNFR2 and CD40 that mediates activation of NF-kappa-B. Rothe et al. (1996) identified ITRAF, which binds to TRAF1, TRAF2, and TRAF3, and that when overexpressed inhibits TRAF2-mediated NF-kappa-B activation. They proposed that ITRAF is an inhibitor of TRAF function that regulates TRAF protein activity by sequestering TRAFs in a latent state in the cytoplasm. Kanamori et al. (2002) found that mouse Traf2 interacted directly with T2bp, and they presented evidence that T2BP is involved in TNF-mediated signaling by its interaction with TRAF2.

### **Application Notes**

Optimal dilution of the TRAF2 antibody should be determined by the researcher.

#### **Immunogen**

Amino acids RLDQDKIEALSSKVQQLERSIGLKDLAMADLE of human TRAF2 were used as the immunogen for the TRAF2 antibody.

#### **Storage**

After reconstitution, the TRAF2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.