

TRAIL Antibody (R32840)


Catalog No.	Formulation	Size
R32840	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug




Citations (1)

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA, 0.025% sodium azide
UniProt	P50591
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This TRAIL antibody is available for research use only.




IHC staining of FFPE human tonsil with TRAIL antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE mouse spleen with TRAIL antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



IHC staining of FFPE rat liver with TRAIL antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.



Western blot testing of human 1) HL-60 and 2) ThP1 cell lysate with TRAIL antibody. Predicted molecular weight: ~34 kDa.

Description

TNF-related apoptosis-inducing ligand (TRAIL), is a protein functioning as a ligand that induces the process of cell death called apoptosis. TRAIL has also been designated CD253 (cluster of differentiation 253). It is a member of Tumor Necrosis Factor Ligand Superfamily. In humans, the gene that encodes for TRAIL is located at chromosome 3q26. TRAIL has got 281 amino acid proteins. TRAIL mRNA is expressed at significant levels in most normal tissues. TRAIL binds to the death receptors DR4 (TRAIL-RI) and DR5 (TRAIL-RII) and the receptors DcR1 and DcR2, leading to activation of specific kinases and transcription of genes.

Application Notes

Optimal dilution of the TRAIL antibody should be determined by the researcher.

Immunogen

A recombinant protein corresponding to amino acids N40-A272 was used as the immunogen for the TRAIL antibody.

Storage

After reconstitution, the TRAIL antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.