


Transthyretin Antibody / TTR / Prealbumin (F54873)


Catalog No.	Formulation	Size
F54873-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54873-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity purified
UniProt	P02766
Localization	Cytoplasmic
Applications	Immunofluorescence : 1:25 Western Blot : 1:1000-1:2000 Immunohistochemistry (FFPE) : 1:25 Flow Cytometry : 1:25 (1x10 ⁶ cells)
Limitations	This Transthyretin antibody is available for research use only.




IHC testing of FFPE human liver tissue with Transthyretin antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Western blot testing of human 1) placenta and 2) plasma lysate with Transthyretin antibody. Predicted molecular weight ~16 kDa.



Immunofluorescent staining of human pancreas and kidney tissue with Transthyretin antibody (green) and DAPI nuclear stain (blue).



Flow cytometry testing of fixed and permeabilized human HepG2 cells with Transthyretin antibody; Blue=isotype control, Green= Transthyretin antibody.

Description

Transthyretin, one of the three prealbumins including alpha-1-antitrypsin, transthyretin and orosomucoid. Transthyretin is a carrier protein; it transports thyroid hormones in the plasma and cerebrospinal fluid, and also transports retinol (vitamin A) in the plasma. The protein consists of a tetramer of identical subunits. More than 80 different mutations in this gene have been reported; most mutations are related to amyloid deposition, affecting predominantly peripheral nerve and/or the heart, and a small portion of the gene mutations is non-amyloidogenic. The diseases caused by mutations include amyloidotic polyneuropathy, euthyroid hyperthyroxinaemia, amyloidotic vitreous opacities, cardiomyopathy, oculoleptomeningeal amyloidosis, meningocerebrovascular amyloidosis, carpal tunnel syndrome, etc.

Application Notes

The stated application concentrations are suggested starting points. Titration of the Transthyretin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 71-98 from the human protein was used as the immunogen for the Transthyretin antibody.

Storage

Aliquot the Transthyretin antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

