


TRAP1 Antibody / HSP75 (R32204)


Catalog No.	Formulation	Size
R32204	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	Q12931
Localization	Cytoplasmic
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This TRAP1 antibody is available for research use only.



Western blot testing of 1) rat heart, 2) rat kidney, 3) rat brain, 4) human SMMC, 5) human PANC and 6) human A549 lysate with TRAP1 antibody. Expected molecular weight: 75-80 kDa.



IHC testing of FFPE human intestinal cancer tissue with TRAP1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

Description

Heat shock protein 75 kDa, mitochondrial is a protein that in humans is encoded by the TRAP1 gene. It is mapped to 16p13.3. This gene encodes a mitochondrial chaperone protein that is member of the heat shock protein 90 (HSP90) family. The encoded protein has ATPase activity and interacts with tumor necrosis factor type I. And this protein may function in regulating cellular stress responses. In addition, it was found that TRAP1 interacted with the N-terminal half of TNFR1. Also, TRAP1 interacted with the C-terminal ends of the proteins encoded by both multiple exostoses-causing genes, EXT1 and EXT2, but not with EXT1 or EXT3.

Application Notes

Optimal dilution of the TRAP1 antibody should be determined by the researcher.

Immunogen

Amino acids 571-704 of human TRAP1 were used as the immunogen for the TRAP1 antibody.

Storage

After reconstitution, the TRAP1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.