

## TRAPPC2 Antibody (F40039)

Catalog No.	Formulation	Size
F40039-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F40039-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Predicted Reactivity</b>	Mouse, Bovine, Pig, Chicken, Zebrafish
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	P0DI81
<b>Applications</b>	Western Blot : 1:1000
<b>Limitations</b>	This TRAPPC2 antibody is available for research use only.



TRAPPC2 antibody western blot analysis in NCI-H292 lysate.

### Description

TRAPPC2 is thought to be part of a large multi-subunit complex involved in the targeting and fusion of endoplasmic reticulum-to-Golgi transport vesicles with their acceptor compartment. In addition, the encoded protein can bind c-myc promoter-binding protein 1 and block its transcriptional repression capability. Mutations in this gene are a cause of spondyloepiphyseal dysplasia tarda (SEDT). A processed pseudogene of this gene is located on chromosome 19, and other pseudogenes are found on chromosomes 8 and Y.

## Application Notes

Titration of the TRAPPC2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 11-39 from the human protein was used as the immunogen for this TRAPPC2 antibody.

## Storage

Aliquot the TRAPPC2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.