



TRF2 Antibody / Telomeric repeat-binding factor 2 (R32769)

Catalog No.	Formulation	Size
R32769	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose, 0.025% sodium azide
UniProt	Q15554
Localization	Nuclear
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This TRF2 antibody is available for research use only.





IHC testing of FFPE human tonsil tissue with TRF2 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



IHC testing of FFPE human lung cancer tissue with TRF2 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.

Description

Telomeric repeat-binding factor 2, also known as TERF2, TRF2, or TRBF2, is encoded in humans by the TERF2 gene. This gene encodes a telomere specific protein, TERF2, which is a component of the telomere nucleoprotein complex. This protein is present at telomeres in metaphase of the cell cycle, is a second negative regulator of telomere length and plays a key role in the protective activity of telomeres. While having similar telomere binding activity and domain organization, TERF2 differs from TERF1 in that its N terminus is basic rather than acidic.

Application Notes

Optimal dilution of the TRF2 antibody should be determined by the researcher.

Immunogen

Amino acids A81-K287 from the human protein were used as the immunogen for the TRF2 antibody.

Storage

After reconstitution, the TRF2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.