


TRIB2 Antibody (R32292)

Catalog No.	Formulation	Size
R32292	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	Q92519
Localization	Cytoplasmic, membranous
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This TRIB2 antibody is available for research use only.


kDa 200 –	Western blot testing of human SW620 cell lysate with TRIB2 antibody.
116 97	Expected/observed molecular weight ~39 kDa.
66 44	
31	
22 14	
6	



IHC testing of FFPE human tonsil with TRIB2 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE mouse spleen with TRIB2 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat spleen with TRIB2 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

Description

Tribbles homolog 2 is a protein that in humans is encoded by the TRIB2 gene. This gene encodes one of three members of the Tribbles family. The Tribbles members share a Trb domain, which is homologous to protein serine-threonine kinases, but lacks the active site lysine and probably lacks a catalytic function. The Tribbles proteins interact and modulate the activity of signal transduction pathways in a number of physiological and pathological processes. This Tribbles member induces apoptosis of cells mainly of the hematopoietic origin. It has been identified as a protein upregulated by inflammatory stimuli in myeloid (THP-1) cells, and also as an oncogene that inactivates the transcription factor C/EBP alpha (CCAAT/enhancer-binding protein alpha) and causes acute myelogenous leukemia. Alternatively spliced transcript variants have been found for this gene.

Application Notes

Optimal dilution of the TRIB2 antibody should be determined by the researcher.

Immunogen

Amino acids DLKLRKFIFKDEERTRVKLESLEDAYILRGDDDSLSD of human TRIB2 were used as the immunogen for the TRIB2 antibody.

Storage

After reconstitution, the TRIB2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.