
TRIM28 Antibody / KAP1 [clone 9E3.] (RQ5933)


Catalog No.	Formulation	Size
RQ5933	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a
Clone Name	9E3.
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	Q13263
Localization	Nuclear
Applications	Western blot: 0.5-1ug/ml Immunohistochemistry (FFPE and Frozen): 1-2ug/ml Immunofluorescence: 2-4ug/ml Flow cytometry: 1-3ug/million cells
Limitations	This TRIM28 antibody is available for research use only.




IHC staining of FFPE human breast cancer with TRIM28 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human rectal cancer with TRIM28 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat intestine with TRIM28 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of frozen human placenta with TRIM28 antibody.




Immunofluorescent staining of FFPE human rectal cancer with TRIM28 antibody (green). HIER: steam section in pH8 EDTA for 20 min.



Immunofluorescent staining of FFPE human U-2 OS cells with TRIM28 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of human 1) HeLa, 2) PC-3, 3) HEK293, 4) A549, 5) Jurkat, 6) ThP-1, 7) rat PC-12 and 8) mouse NIH 3T3 lysate with TRIM28 antibody. Observed molecular weight: 88~110 kDa depending on sumoylation.



Flow cytometry testing of human A549 cells with TRIM28 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= TRIM28 antibody.

Description

Tripartite motif-containing 28 (TRIM28), also known as transcriptional intermediary factor 1 beta (TIF1b) and KAP1 (KRAB-associated protein-1), is a protein that in humans is encoded by the TRIM28 gene. The protein encoded by this gene mediates transcriptional control by interaction with the Kruppel-associated box repression domain found in many transcription factors. The protein localizes to the nucleus and is thought to associate with specific chromatin regions. KAP1 is a ubiquitously expressed protein involved in many critical functions including: transcriptional regulation, cellular differentiation and proliferation, DNA damage repair, viral suppression, and apoptosis. Its functionality is dependent upon post-translational modifications. Phosphorylation of KAP1 acts as a deactivator of the protein in many of its mechanisms while sumoylation acts as an activator.

Application Notes

Optimal dilution of the TRIM28 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids A699-P835) was used as the immunogen for the TRIM28 antibody.

Storage

After reconstitution, the TRIM28 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.