


TRIM29 Antibody [clone TRIM29/1041] (V2507)


| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V2507-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V2507-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V2507SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |
| V2507IHC-7ML | Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only* | 7 ml |

[Bulk quote request](#)

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG2a, kappa |
| Clone Name | TRIM29/1041 |
| Purity | Protein G affinity chromatography |
| UniProt | Q14134 |
| Localization | Cytoplasmic & cell surface |
| Applications | ELISA : order BSA/sodium azide-free format for coating Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT |
| Limitations | This TRIM29 antibody is available for research use only. |



IHC: Formalin-fixed, paraffin-embedded human lung squamous cell carcinoma stained with TRIM29 antibody (clone TRIM29/1041).



IHC: Formalin-fixed, paraffin-embedded human esophageal carcinoma stained with TRIM29 antibody (clone TRIM29/1041).

Description

It recognizes a 66kDa protein, which is identified as Tripartite motif-containing protein 29 (TRIM29). It interacts with the intermediate filament protein vimentin, a substrate for the PKC family of protein kinases, and with hPKCI-1, an inhibitor of the PKCs. TRIM29 protein contains both zinc finger and leucine zipper motifs, suggesting that it may form homodimers and possibly associate with DNA. High expression of TRIM29 has been reported in gastric cancer and pancreatic cancer, and correlates with enhanced tumor growth and lymph node metastasis. TRIM29 is also able to distinguish lung squamous cell carcinoma from lung adenocarcinoma with ~90% positive accuracy, when used in a panel with TTF-1, p63, CK5/6, and Napsin-A antibodies.

Application Notes

Optimal dilution of the TRIM29 antibody should be determined by the researcher.

1. Staining of formalin/paraffin tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

A recombinant fragment (126 amino acid residues between aa 1-200) of human TRIM29 protein was used as the immunogen for the TRIM29 antibody.

Storage

Store the TRIM29 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).