



TRPC6 Antibody (R30847)

Catalog No.	Formulation	Size
R30847	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug


Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA, 0.025% sodium azide/thimerosal
UniProt	Q9Y210
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml
Limitations	This TRPC6 antibody is available for research use only.





IHC-P: TRPC6 antibody testing of human intestinal cancer tissue





Description

Transient receptor potential cation channel, subfamily C, member 6, also known as TRP6, is a transient receptor potentialion channel. Northern blot analysis revealed that TRPC6 is expressed primarily in placenta, lung, spleen, ovary, and small intestine. The protein is a nonselective cation channel that is activated by diacylglycerol (DAG) in a membrane-delimited fashion, independently of protein kinase C. Although TRPC3, the closest structural relative of TRPC6, is activated in the same manner, human TRPC1 and mouse Trpc4 and Trpc5 were unresponsive to DAG. Immunofluorescence studies showed that most TRPC6 expression is confined to podocytes. It is also expressed in glomerular endothelial cells. Cardiac-specific overexpression of the protein in transgenic mice resulted in heightened sensitivity to stress, a propensity for lethal cardiac growth and heart failure, and an increase in Nfat-dependent expression of beta-myosin heavy chain(MYH7), a marker for pathologic hypertrophy. Studies of Trpc6 -/- mice showed that it has a role in regulation of smooth muscle tone in blood vessels and lung.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the TRPC6 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the middle region of human TRPC6 (HDYFCKCNDCNQKQKHD) was used as the immunogen for this TRPC6 antibody.

Storage

After reconstitution, the TRPC6 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.