


TRPM4 Antibody (R32286)


Catalog No.	Formulation	Size
R32286	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q8TD43
Localization	Cytoplasmic, membranous
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This TRPM4 antibody is available for research use only.



IHC staining of FFPE human prostate cancer tissue with TRPM4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human squamous cell carcinoma of cervix tissue with TRPM4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Description

Transient receptor potential cation channel subfamily M member 4 (hTRPM4), also known as melastatin-4, is a protein that in humans is encoded by the TRPM4 gene. It is mapped to 19q13.33. The protein encoded by this gene is a calcium-activated nonselective ion channel that mediates transport of monovalent cations across membranes, thereby depolarizing the membrane. The activity of the encoded protein increases with increasing intracellular calcium concentration, but this channel does not transport calcium. Two transcript variants encoding different isoforms have been found for this gene.

Application Notes

Optimal dilution of the TRPM4 antibody should be determined by the researcher.

Immunogen

Amino acids 1079-1214 of human TRPM4 were used as the immunogen for the TRPM4 antibody.

Storage

After reconstitution, the TRPM4 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.