


TSG6 Antibody (R32752)

| Catalog No. | Formulation | Size |
|-------------|---|--------|
| R32752 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

Bulk quote request

| Availability | 1-3 business days |
|--------------------|--|
| Species Reactivity | Human, Rat |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit IgG |
| Purity | Antigen affinity |
| Buffer | Lyophilized from 1X PBS with 2.5% BSA, 0.025% sodium azide |
| UniProt | P98066 |
| Applications | Western Blot : 0.5-1ug/ml |
| Limitations | This TSG6 antibody is available for research use only. |



Description

Tumor necrosis factor-inducible gene 6 protein, also known as TSG-6, is a protein that in humans is encoded by the TNFAIP6 gene. The protein encoded by this gene is a secretory protein that contains a hyaluronan-binding domain, and thus is a member of the hyaluronan-binding protein family. The hyaluronan-binding domain is known to be involved in extracellular matrix stability and cell migration. This protein has been shown to form a stable complex with inter-alpha-inhibitor (I alpha I), and thus enhance the serine protease inhibitory activity of I alpha I, which is important in the protease network associated with inflammation. This gene can be induced by proinflammatory cytokines such as tumor necrosis factor alpha and interleukin-1. Enhanced levels of this protein are found in the synovial fluid of patients with osteoarthritis and rheumatoid arthritis.

Application Notes

Optimal dilution of the TSG6 antibody should be determined by the researcher.

Immunogen

Amino acids 46-91 (KYKLTYAEAKAVCEFEGGHLATYKQLEAARKIGFHVCAAGWMAKGR) from the human protein were used as the immunogen for the TSG6 antibody.

Storage

After reconstitution, the TSG6 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.