


TTF-1 Antibody / NKX2.1 [clone TTF1/8430] (V4344)

Catalog No.	Formulation	Size
V4344-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4344-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4344SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	TTF1/8430
Purity	Protein A/G affinity
UniProt	P43699
Localization	Nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This TTF-1 antibody is available for research use only.



IHC staining of FFPE human lung adenocarcinoma tissue with TTF-1 antibody (clone TTF1/8430). Inset: PBS used in place of primary Ab (secondary Ab negative control).
 HIERS: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

SDS-PAGE analysis of purified, BSA-free TTF-1 antibody (clone TTF1/8430) as confirmation of integrity and purity.



Description

Recognizes a protein of 40kDa, identified as Thyroid transcription factor-1 (TTF-1). TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver, stomach, pancreas, small intestine, colon, kidney, breast, skin, testes, pituitary, prostate, and adrenal glands are unreactive. Anti-TTF-1 is useful in differentiating primary adenocarcinoma of the lung from metastatic carcinomas originating in the breast, mediastinal germ cell tumors, and malignant mesothelioma. It can also be used to differentiate small cell lung carcinoma from lymphoid infiltrates. Loss of TTF-1 expression in non-small cell lung carcinoma has been associated with aggressive behavior of such neoplasms. TTF-1 reactivity is also seen in thyroid malignancies.

Application Notes

Optimal dilution of the TTF-1 antibody should be determined by the researcher.

Immunogen

Recombinant human TTF-1 protein was used as the immunogen for the TTF-1 antibody.

Storage

Aliquot the TTF-1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.