


UchL1 Antibody [clone UCHL1/841] (V2921)


Catalog No.	Formulation	Size
V2921-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2921-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2921SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	UCHL1/841
Purity	Protein G affinity chromatography
UniProt	P09936
Localization	Cytoplasmic, ER membrane
Applications	Western Blot : 1-2ug/ml
Limitations	This UchL1 antibody is available for research use only.



Western blot testing of human brain lysate with UchL1 antibody (clone UCHL1/841). Predicted molecular weight ~25 kDa.



Description

This mAb reacts with a protein of 20-30kDa, identified as PGP9.5, also known as ubiquitin carboxyl-terminal hydrolase-1 (UchL1). Initially, PGP9.5 expression in normal tissues was reported in neurons and neuroendocrine cells but later it was found in distal renal tubular epithelium, spermatogonia, Leydig cells, oocytes, melanocytes, prostatic secretory epithelium, ejaculatory duct cells, epididymis, mammary epithelial cells, Merkel cells, and dermal fibroblasts. Furthermore, immunostaining for PGP9.5 has been shown in a wide variety of mesenchymal neoplasms as well. A mutation in PGP9.5 gene is believed to cause a form of Parkinson's disease.

Application Notes

Optimal dilution of the UchL1 antibody should be determined by the researcher.

Immunogen

Recombinant human protein was used as the immunogen for the UchL1 antibody.

Storage

Store the UchL1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).