


UPF3B Antibody / RENT3B (R32318)


Catalog No.	Formulation	Size
R32318	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	Q9BZI7
Localization	Cytoplasmic
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This UPF3B antibody is available for research use only.




Western blot testing of 1) rat heart, 2) rat brain, and human 3) HeLa, 4) 22RV1, 5) U87 and 6) Jurkat lysate with UPF3B antibody. Expected/observed molecular weight ~58 kDa.



IHC testing of FFPE human placenta with UPF3B antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE mouse testis with UPF3B antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat testis with UPF3B antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

Description

Regulator of nonsense transcripts 3B is a protein that in humans is encoded by the UPF3B gene. This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. The encoded protein is one of two functional homologs to yeast Upf3p. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein binds to the mRNA and remains bound after nuclear export, acting as a nucleocytoplasmic shuttling protein. It forms with Y14 a complex that binds specifically 20 nt upstream of exon-exon junctions. This gene is located on the long arm of chromosome X. Two splice variants encoding different isoforms have been found for this gene.

Application Notes

Optimal dilution of the UPF3B antibody should be determined by the researcher.

Immunogen

Amino acids SEKTEKKEEVVKRDRIRNKDRPAMQLYQPGARSRNRL of human UPF3B were used as the immunogen for the UPF3B antibody.

Storage

After reconstitution, the UPF3B antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.