


Uroplakin 1A Antibody Microarray Validated / UPK1A [clone UPK1A/2925] (V7937)


Catalog No.	Formulation	Size
V7937-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7937-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7937SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	UPK1A/2925
Purity	Protein G affinity chromatography
UniProt	O00322
Localization	Cell surface, cytoplasmic
Applications	ELISA (order BSA-free Format For Coating) : Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Uroplakin 1A antibody is available for research use only.




Immunohistochemistry of Uroplakin 1A antibody in human urothelial carcinoma. Formalin-fixed, paraffin-embedded human urothelial carcinoma tissue was stained with microarray validated Uroplakin 1A antibody (clone UPK1A/2925) following heat-induced epitope retrieval by boiling in pH 9 10 mM Tris with 1 mM EDTA for 10-20 minutes and cooling prior to testing. HRP-DAB brown chromogenic signal demonstrates strong membranous staining in tumor cells, consistent with apical membrane localization and urothelial differentiation associated with Uroplakin 1A, while surrounding stromal cells show minimal background staining.




Immunohistochemistry of Uroplakin 1A antibody in human urothelial carcinoma. Formalin-fixed, paraffin-embedded human urothelial carcinoma tissue was stained with microarray validated Uroplakin 1A antibody (clone UPK1A/2925) following heat-induced epitope retrieval by boiling in pH 9 10 mM Tris with 1 mM EDTA for 10-20 minutes and cooling prior to testing. HRP-DAB brown chromogenic signal highlights prominent membranous staining in malignant urothelial cells, consistent with the apical membrane localization and differentiation-associated expression pattern of Uroplakin 1A, while adjacent stromal elements remain largely negative.

Human Protein Microarray Specificity Validation



HuProt protein microarray specificity analysis of Uroplakin 1A antibody (clone UPK1A/2925). The antibody was evaluated against a HuProt array containing more than 19,000 full-length human proteins. Signal intensity is expressed as a Z-score, representing the number of standard deviations above the mean signal generated across the array when the antibody, together with a fluorescently labeled anti-IgG secondary antibody, binds to a given protein. Proteins are ranked in descending order by Z-score, and the S-score reflects the difference in Z-scores between the top-ranked protein and the next highest signal, providing a quantitative measure of relative target specificity. Clone UPK1A/2925 demonstrates a markedly elevated Z-score for Uroplakin 1A with strong S-score separation from other proteins, supporting highly specific recognition of UPK1A on the array and confirming its microarray validated status.



SDS-PAGE analysis of purified, BSA-free Uroplakin 1A antibody (clone UPK1A/2925) as confirmation of integrity and purity.

Description

Uroplakin 1A antibody recognizes Uroplakin 1A, a urothelial membrane protein encoded by the UPK1A gene and a critical structural component of the apical surface of umbrella cells. Uroplakin 1A Antibody Microarray Validated (clone UPK1A/2925) is a mouse monoclonal antibody developed for research applications requiring specific detection of this differentiation-associated protein in urothelial tissues. Uroplakin 1A localizes predominantly to the apical plasma membrane of bladder, ureter, and renal pelvis umbrella cells, where it contributes to formation of specialized urothelial plaques that maintain epithelial barrier function and polarity.

Uroplakin 1A antibody, also referred to as UPK1A antibody and UPlA antibody, targets a member of the uroplakin family of integral membrane proteins. Uroplakin 1A contains multiple transmembrane domains and extracellular loops that participate in heterodimer formation with Uroplakin 2. These heterodimers assemble into asymmetric unit membrane plaques that provide mechanical stability and reduce permeability of the urothelial surface to urine and solutes. This highly specialized membrane architecture protects underlying tissues from osmotic stress and toxic urinary metabolites.

UPK1A expression is tightly restricted to terminally differentiated urothelial umbrella cells and is closely associated with epithelial maturation and maintenance of apical membrane specialization. Because of this tissue-specific distribution, Uroplakin 1A is widely used as a marker of urothelial lineage in normal tissue biology and tumor classification research. Membranous staining patterns are particularly informative in immunohistochemical analyses of bladder and upper urinary tract specimens.

In oncology research, UPK1A expression is frequently evaluated in urothelial carcinoma and in the differential diagnosis of metastatic carcinomas of unknown primary origin. Detection of Uroplakin 1A supports urothelial differentiation, whereas

most non-urothelial epithelial tumors lack expression. This selective expression profile makes Uroplakin 1A antibody valuable for studies of bladder cancer biology and epithelial lineage determination.

Microarray validation of clone UPK1A/2925 supports its specificity across large-scale human protein panels, providing confidence in target recognition for tissue-based and proteomic research applications at NSJ Bioreagents.

Application Notes

Optimal dilution of the Uroplakin 1A antibody should be determined by the researcher.

Immunogen

Amino acids 114-173 were used as the immunogen for the Uroplakin 1A antibody.

Storage

Store the Uroplakin 1A antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).