


Vcam-1 Antibody (R32890)


Catalog No.	Formulation	Size
R32890	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P29533
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml ELISA (Capture; Recombinant Mouse Protein) : 0.1-0.5ug/ml
Limitations	This Vcam-1 antibody is available for research use only.



Western blot testing of 1) mouse spleen, 2) mouse lung, 3) rat spleen and 4) rat lung with Vcam-1 antibody at 0.5ug/ml. Expected molecular weight: 74-80 kDa (unmodified), 100-110 kDa (glycosylated).



IHC testing of FFPE mouse spleen tissue with Vcam-1 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



IHC testing of FFPE rat spleen tissue with Vcam-1 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.

Description

CD106 (cluster of differentiation 106) also known as vascular cell adhesion molecule 1 (VCAM-1), is a protein that in humans is encoded by the VCAM1 gene. In inflammatory conditions and in cardiac allografts undergoing rejection, VCAM1 is upregulated in endothelium of postcapillary venules. Arterial expression of VCAM1 is also found in experimental models of atherosclerosis in the rabbit. This gene is mapped to chromosome 1 by Southern analysis of somatic cell hybrids. VCAM-1 functions as a cell adhesion molecule. The VCAM-1 protein mediates the adhesion of lymphocytes, monocytes, eosinophils, and basophils to vascular endothelium. It also functions in leukocyte-endothelial cell signal transduction, and it may play a role in the development of atherosclerosis and rheumatoid arthritis. CAM741 works similar to cotransin in that it represses the biosynthesis of VCAM1 cells by blocking the process of cotranslational translocation, which is dependent on the signal peptide of VCAM1. Among the lung metastasis signature genes identified, several, including VCAM1, were functionally validated.

Application Notes

Optimal dilution of the Vcam-1 antibody should be determined by the researcher.

Immunogen

A recombinant mouse protein corresponding to amino acids F25-L270 was used as the immunogen for the Vcam-1 antibody.

Storage

After reconstitution, the Vcam-1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.