


Vimentin Antibody [clone LN-6] (V2927)

Catalog No.	Formulation	Size
V2927-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2927-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2927SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2927IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Citations (11)

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgM, kappa
Clone Name	LN-6
Purity	PEG precipitation
UniProt	P08670
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE): 0.1-0.2ug/ml for 30 min at RT
Limitations	This Vimentin antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human melanoma stained with Vimentin antibody (clone LN-6).

Description

This mAb reacts with a 58kDa protein identified as vimentin. It reacts with a non-hematopoietic epitope of vimentin and shows no cross-reaction with other closely related intermediate filament proteins (IFP's) such as desmin, keratin, neurofilament, and glial fibrillary acid protein. Vimentin is ubiquitously expressed in mesenchymal cells such as fibroblasts, smooth muscle cells, and endothelium. Antibody against vimentin is useful as part of an antibody panel for differential diagnosis of tumors of unknown origin. Ab-2 does not react with leukocyte common antigen-positive tissues such as lymphomas and leukemias.

Application Notes

Optimal dilution of the Vimentin antibody should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

A human thymic nuclear extract was used as the immunogen for the Vimentin antibody.

Storage

Store the Vimentin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).