


# **VLDL Receptor Antibody / VLDLR (RQ6695)**


| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| RQ6695      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

## **Bulk quote request**

| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human, Mouse, Rat  |
| Format             | Antigen affinity purified  |
| Clonality          | Polyclonal (rabbit origin)   |
| Isotype            | Rabbit IgG   |
| Purity             | Antigen affinity purified  |
| Buffer             | Lyophilized from 1X PBS with 2% Trehalose  |
| UniProt            | P98155   |
| Applications       | Western Blot : 1-2ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml |
| Limitations        | This VLDL Receptor antibody is available for research use only.                          |



Flow cytometry testing of human JK-1 cells with VLDL Receptor antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= VLDL Receptor antibody.



Western blot testing of 1) human ThP-1, 2) human Jurkat, 3) rat brain, 4) rat C6, 5) mouse heart, 6) mouse brain and 7) mouse Neuro-2a antibody. Expected molecular weight: 96-160 kDa depending on glycosylation level.

#### **Description**

The very-low-density-lipoprotein receptor (VLDLR) is a transmembrane lipoprotein receptor of the low-density-lipoprotein (LDL) receptor family. The low density lipoprotein receptor (LDLR) gene family consists of cell surface proteins involved in receptor-mediated endocytosis of specific ligands. This gene encodes a lipoprotein receptor that is a member of the LDLR family and plays important roles in VLDL-triglyceride metabolism and the reelin signaling pathway. Mutations in this gene cause VLDLR-associated cerebellar hypoplasia. Alternative splicing generates multiple transcript variants encoding distinct isoforms for this gene.

## **Application Notes**

Optimal dilution of the VLDL Receptor antibody should be determined by the researcher.

## **Immunogen**

Recombinant human protein (amino acids R231-S770) was used as the immunogen for the VLDL Receptor antibody.

#### **Storage**

After reconstitution, the VLDL Receptor antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.