

## von Willebrand Factor Antibody [clone F8/86] (V3233)


Catalog No.	Formulation	Size
V3233-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3233-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3233SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug




Citations (23)

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	F8/86
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P04275
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 0.1-0.2ug/ml for 30 min at RT
<b>Limitations</b>	This von Willebrand Factor antibody is available for research use only.



IHC testing of FFPE human tonsil with von Willebrand Factor antibody (clone F8/86). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE Analysis of Purified, BSA-Free von Willebrand Factor Antibody (clone F8/86). Confirmation of Integrity and Purity of the Antibody.

## Description

von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.

## Application Notes

The optimal dilution of the von Willebrand Factor antibody for each application should be determined by the researcher.

## Immunogen

Human plasma vWF protein was used as the immunogen for this von Willebrand Factor antibody.

## Storage

Store the von Willebrand Factor antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).