


von Willebrand Factor Antibody / VWF (RQ7212)


Catalog No.	Formulation	Size
RQ7212	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P04275
Applications	Immunohistochemistry (FFPE) : 2-5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This von Willebrand Factor antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with von Willebrand Factor antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human thyroid cancer tissue with von Willebrand Factor antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human renal cancer tissue with von Willebrand Factor antibody.
HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

Von Willebrand factor (VWF) is a blood glycoprotein involved in hemostasis. It is mapped to 12p13.31. The VWF gene encodes von Willebrand factor (VWF), a large multimeric glycoprotein that plays a central role in the blood coagulation system, serving both as a major mediator of platelet-vessel wall interaction and platelet adhesion, and as a carrier for coagulation factor VIII. VWF released from endothelial cell Weibel-Palade bodies bound particularly avidly to the extracellular matrix. VWF deficiency or dysfunction (von Willebrand disease) leads to a bleeding tendency, which is most apparent in tissues having high blood flow shear in narrow vessels.

Application Notes

Optimal dilution of the von Willebrand Factor antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids R1287-Q2770) was used as the immunogen for the von Willebrand Factor antibody.

Storage

After reconstitution, the von Willebrand Factor antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.