


vWF Antibody / von Willebrand Factor [clone VWF/2480] (V8164)


Catalog No.	Formulation	Size
V8164-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8164-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8164SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	VWF/2480
Purity	Protein G affinity chromatography
UniProt	P04275
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This vWF antibody is available for research use only.



IHC staining of FFPE human tonsil with vWF antibody (clone VWF/2480). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free vWF antibody (clone VWF/2480) as confirmation of integrity and purity.

Description

Von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.

Application Notes

Optimal dilution of the vWF antibody should be determined by the researcher.

Immunogen

A recombinant full-length human protein was used as the immunogen for this vWF antibody.

Storage

Store the vWF antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).