


# WASP Antibody (R32099)

Catalog No.	Formulation	Size
R32099	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug


## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P42768
Localization	Cytoplasmic
Applications	Western Blot: 0.1-0.5ug/ml Immunocytochemistry: 0.5-1ug/ml IHC (Frozen): 0.5-1ug/ml IHC (FFPE): 0.5-1ug/ml
Limitations	This WASP antibody is available for research use only.




Western blot testing of 1) rat liver, 2) human placenta, 3) rat spleen, 4) rat pancreas, 5) human HepG2, 6) HeLa, 7) HEPA and 8) 22RV1 lysate with WASP antibody.


Expected/observed molecular weight ~53 kDa.



IHC testing of FFPE human lung cancer tissue with WASP antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of frozen human placenta with WASP antibody.



ICC testing of A549 cells with WASP antibody.

#### **Description**

The Wiskott-Aldrich syndrome (WAS) family of proteins share similar domain structure, and are involved in transduction of signals from receptors on the cell surface to the actin cytoskeleton. The presence of a number of different motifs suggests that they are regulated by a number of different stimuli, and interact with multiple proteins. Recent studies have demonstrated that these proteins, directly or indirectly, associate with the small GTPase, Cdc42, known to regulate formation of actin filaments, and the cytoskeletal organizing complex, Arp2/3. Wiskott-Aldrich syndrome is a rare, inherited, X-linked, recessive disease characterized by immune dysregulation and microthrombocytopenia, and is caused by mutations in the WAS gene. The WAS gene product is a cytoplasmic protein, expressed exclusively in hematopoietic cells, which show signalling and cytoskeletal abnormalities in WAS patients. A transcript variant arising as a result of alternative promoter usage, and containing a different 5' UTR sequence, has been described, however, its full-length nature is not known.

### **Application Notes**

Optimal dilution of the WASP antibody should be determined by the researcher.

#### **Immunogen**

Amino acids ADEDEAQAFRALVQEKIQKRNQRQSGDR of human WASP were used as the immunogen for the WASP antibody.

#### **Storage**

After reconstitution, the WASP antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.