


## WNT2B Antibody (R32298)


Catalog No.	Formulation	Size
R32298	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
<b>UniProt</b>	Q93097
<b>Applications</b>	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
<b>Limitations</b>	This WNT2B antibody is available for research use only.




Western blot testing of human 22RV1 cell lysate with WNT2B antibody.  
Expected/observed molecular weight ~44 kDa.



IHC testing of FFPE human lung cancer tissue with WNT2B antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE mouse brain with WNT2B antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat brain with WNT2B antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

## Description

WNT2B (Wingless-Type MMTV Integration Site Family Member 2B), is a protein that in humans is encoded by the WNT2B gene. This gene encodes a member of the wingless-type MMTV integration site (WNT) family of highly conserved, secreted signaling factors. Katoh et al. (1996) used fluorescence in situ hybridization to map the WNT13 gene to chromosome 1p13. WNT family members function in a variety of developmental processes including regulation of cell growth and differentiation and are characterized by a WNT-core domain. This gene may play a role in human development as well as human carcinogenesis.

## Application Notes

Optimal dilution of the WNT2B antibody should be determined by the researcher.

## Immunogen

Amino acids QRYPDIMRSVGEAREWIRECQHQR of human WNT2B were used as the immunogen for the WNT2B antibody.

## Storage

After reconstitution, the WNT2B antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.