



WT1 Antibody (Wilms Tumor protein) (F41819)

Catalog No.	Formulation	Size
F41819-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F41819-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Mouse, Pig, Rat, Xenopus
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	P19544
Localization	Nuclear, cytoplasmic
Applications	Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 Immunofluorescence : 1:10-1:50 Flow Cytometry : 1:10-1:50
Limitations	This WT1 antibody is available for research use only.






WT1 antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue.



Confocal immunofluorescent analysis of WT1 antibody with MCF-7 cells followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green). DAPI was used to stain the nuclei (blue).



WT1 antibody flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control (left histogram). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.

Description

Wilms Tumor protein is a transcription factor that contains four zinc-finger motifs at the C-terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with Wilm's tumors. This gene exhibits complex tissue-specific and polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal and paternal alleles in different tissues. Multiple transcript variants have been described. In several variants, there is evidence for the use of a non-AUG (CUG) translation initiation site upstream of and in-frame with the first AUG. Authors of PMID:7926762 also provide evidence that WT1 mRNA undergoes RNA editing in human and rat, and that this process is tissue-restricted and developmentally regulated. [provided by RefSeq].

Application Notes

Titration of the WT1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 346-375 from the human protein was used as the immunogen for this WT1 antibody.

Storage

Aliquot the WT1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.