


## XIAP associated factor 1 Antibody / XAF1 (RQ7213)


Catalog No.	Formulation	Size
RQ7213	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	Q6GPH4
<b>Localization</b>	Cytoplasmic, nuclear
<b>Applications</b>	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
<b>Limitations</b>	This XIAP associated factor 1 antibody is available for research use only.




IHC staining of FFPE human thyroid cancer tissue with XIAP associated factor 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placental tissue with XIAP associated factor 1 antibody.  
HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human lung cancer tissue with XIAP associated factor 1 antibody.  
HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) placenta, 2) 293T and 3) Jurkat cell lysate with XIAP associated factor 1 antibody. Predicted molecular weight ~34 kDa.



Flow cytometry testing of human U-87 MG cells with XIAP associated factor 1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= XIAP associated factor 1 antibody.

## Description

XAF1, also called XIAP-associated factor 1, antagonizes the anti-caspase activity of XIAP and may be important in mediating apoptosis resistance in cancer cells. By genomic sequence analysis, the XAF1 gene contains 7 exons spanning 18 kb. Southern blot analysis suggested that XAF1 is a single-copy gene. The XAF1 gene is mapped on 17p13.1. Fluorescence microscopy demonstrated nuclear expression of XAF1, in contrast to the cytoplasmic expression of XIAP. Functional analysis indicated that XAF1 reverses the XIAP-mediated inhibition of CASP3 and its protection against apoptosis. Northern blot analysis detected wide expression of 3.9-, 4.5-, 6.0-, and 7.0-kb XAF1 transcripts, with highest levels in heart and ovary and lowest levels in brain and testis. Antisense experiments determined that loss of XAF1 expression enhances resistance to apoptosis. Alteration in XAF1 and XIAP RNA expression levels may lead to increased apoptotic resistance and proliferation due to unregulated XIAP function in cancer cells.

## Application Notes

Optimal dilution of the XIAP associated factor 1 antibody should be determined by the researcher.

## Immunogen

Recombinant human protein (amino acids Q197-S301) was used as the immunogen for the XIAP associated factor 1 antibody.

## Storage

After reconstitution, the XIAP associated factor 1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.