


# ZAP70 Antibody [clone 9D5] (RQ5941)


Catalog No.	Formulation	Size
RQ5941	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	9D5
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P43403
Applications	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This ZAP70 antibody is available for research use only.



Flow cytometry testing of human Jurkat cells with ZAP70 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= ZAP70 antibody.



Flow cytometry testing of human 293T cells with ZAP70 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= ZAP70 antibody.



Western blot testing of 1) human Jurkat, 2) human CCRF-CEM, 3) rat thymus, 4) mouse spleen and 5) mouse thymus lysate with ZAP70 antibody. Expected molecular weight ~70 kDa.

### **Description**

ZAP-70 (Zeta-chain-associated protein kinase 70) encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.

#### **Application Notes**

Optimal dilution of the ZAP70 antibody should be determined by the researcher.

#### **Immunogen**

Amino acids MRKKQIDVAIKVLKQGTEKADTEEMMREAQIMHQL from the human protein were used as the immunogen for the ZAP70 antibody.

#### **Storage**

After reconstitution, the ZAP70 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.