


ZBTB7C Antibody / KR-POK [clone PCRP-ZBTB7C-4E12] (V9227)


Catalog No.	Formulation	Size
V9227-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9227-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9227SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	PCRP-ZBTB7C-4E12
Purity	Protein A/G affinity
UniProt	A1YPR0
Localization	Nucleus
Applications	Flow Cytometry : 1-2ug/million cells Immunofluorescence : 1-2ug/ml
Limitations	This ZBTB7C antibody is available for research use only.



Immunofluorescent staining of PFA-fixed human HeLa cells using ZBTB7C antibody (green, clone PCRP-ZBTB7C-4E12) and phalloidin (red).




FACS staining of PFA-fixed human HeLa cells with ZBTB7C antibody (blue, clone PCRP-ZBTB7C-4E12) and isotype control (red).

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using ZBTB7C antibody (clone PCRP-ZBTB7C-4E12). These results demonstrate the foremost specificity of the PCRP-ZBTB7C-4E12 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free ZBTB7C antibody (PCRP-ZBTB7C-4E12) as confirmation of integrity and purity.

Description

Zinc finger and BTB domain-containing protein 7C (ZBTB7C) is involved in negative regulation of cell population proliferation.

Application Notes

Optimal dilution of the ZBTB7C antibody should be determined by the researcher.

Immunogen

Recombinant full-length human ZBTB7C protein was used as the immunogen for the ZBTB7C antibody.

Storage

Aliquot the ZBTB7C antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

