


# Zebrafish Hsc70 Antibody / Hspa8 (RZ1113)


Catalog No.	Formulation	Size
RZ1113	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**


Availability	2-3 weeks
Species Reactivity	Zebrafish
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity chromatography
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q90473
Localization	Cytoplasm, cell membrane, nucleolus
Applications	Western Blot : 0.5-1 ug/ml Immunohistochemistry (FFPE) : 2-5 ug/ml
Limitations	This Zebrafish Hsc70 antibody is available for research use only.



IHC staining of FFPE zebrafish brain tissue with Zebrafish Hsc70 antibody, HRP secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot analysis of Hsc70/HSPA8 protein using Hsc70/HSPA8 antibody and 1) zebrafish head, 2) female zebrafish viscera, 3) male zebrafish viscera, 4) whole female zebrafish, 5) whole male zebrafish and 6) zebrafish embryo tissue lysate. Predicted molecular weight ~71 kDa.



IHC staining of FFPE zebrafish retina tissue with Zebrafish Hsc70 antibody, HRP secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

### **Description**

HSPA8 (heat shock 70kDa protein 8) also known as HSC70, HSC71, HSP73, HSPA10, FORMERLY, LAP1 or LPS-ASSOCIATED PROTEIN 1, is a heat shock protein that in humans is encoded by the HSPA8 gene. The HSPA8 gene contains 9 exons and spans 5 kb. The deduced HSPA8 protein has 646 amino acids and a predicted molecular mass of 70,899 Da. And the HSPA8 gene is mapped on 11q24.1. HSPA8 plays an important role in cells by transiently associating with nascent polypeptides to facilitate correct folding. HSP73 also functions as an ATPase in the disassembly of clathrin-coated vesicles during transport of membrane components through the cell. Rapid decay involves AU-rich binding protein AUF1, which complexes with heat-shock proteins HSC70 and HSP70, translation initiation factor EIF4G, and poly (A)-binding protein. In the absence of II3, Hsc70 formed a complex with Hsp40 and Hip, and this complex, in association with Eif4g and Pabp, formed a high-stability complex with Bim mRNA that protected it from ribonucleases.

#### **Application Notes**

Optimal dilution of the Zebrafish Hsc70 antibody should be determined by the researcher.

#### **Immunogen**

An E.coli-derived zebrafish Hsc70/HSPA8 recombinant protein (amino acids K539-K601) was used as the immunogen for the Zebrafish Hsc70 antibody.

#### **Storage**

After reconstitution, the Zebrafish Hsc70 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.