


ZNF846 Antibody [clone ZNF846/2687] (V8492)

Catalog No.	Formulation	Size
V8492-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8492-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8492SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	ZNF846/2687
Purity	Protein G affinity chromatography
UniProt	Q147U1
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This ZNF846 antibody is available for research use only.




IHC staining of FFPE human breast carcinoma with ZNF846 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using ZNF846 antibody. These results demonstrate the foremost specificity of the ZNF846/2687 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free ZNF846 antibody as confirmation of integrity and purity.

Description

ZNF846 Antibody detects Zinc Finger 846 protein. Zinc finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc finger proteins contain a Kruppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF846 may be involved in transcriptional regulation.

This antibody can be compared with our [ZNF846 Antibody / Zinc Finger Transcription Regulator Antibody](#) (clone PCRP-ZNF846-1E12) for consistent detection of ZNF846 in nuclear and transcriptional regulation studies.

Application Notes

Optimal dilution of the ZNF846 antibody should be determined by the researcher.

Immunogen

Recombinant full-length human ZNF846 protein was used as the immunogen for the ZNF846 antibody.

Storage

Store the ZNF846 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).